

MEDICATION GUIDE

LAMICTAL[®] (la-MIK-tal) XR[™] (lamotrigine) Extended-Release Tablets

Read this Medication Guide before you start taking LAMICTAL XR and each time you get a refill. There may be new information. This information does not take the place of talking with your healthcare provider about your medical condition or treatment. If you have questions about LAMICTAL XR, ask your healthcare provider or pharmacist.

What is the most important information I should know about LAMICTAL XR?

1. LAMICTAL XR may cause a serious skin rash that may cause you to be hospitalized or to stop LAMICTAL XR; it may rarely cause death.

There is no way to tell if a mild rash will develop into a more serious reaction. These serious skin reactions are more likely to happen when you begin taking LAMICTAL XR, within the first 2 to 8 weeks of treatment. But it can happen in people who have taken LAMICTAL XR for any period of time. Children between 2 to 16 years of age have a higher chance of getting this serious skin reaction while taking lamotrigine. LAMICTAL XR is not approved for use in children less than 13 years of age.

The risk of getting a rash is higher if you:

- take LAMICTAL XR while taking valproate [DEPAKENE (valproic acid) or DEPAKOTE (divalproex sodium)].
- take a higher starting dose of LAMICTAL XR than your healthcare provider prescribed.
- increase your dose of LAMICTAL XR faster than prescribed.

LAMICTAL XR can also cause other types of allergic reactions or serious problems that may affect organs and other parts of your body like the liver or blood cells. You may or may not have a rash with these types of reactions.

Call your healthcare provider right away if you have any of the following:

- a skin rash
- hives
- fever
- swollen lymph glands
- painful sores in the mouth or around your eyes

- **swelling of your lips or tongue**
- **yellowing of your skin or eyes**
- **unusual bruising or bleeding**
- **severe fatigue or weakness**
- **severe muscle pain**
- **frequent infections**

These symptoms may be the first signs of a serious reaction. A healthcare provider should examine you to decide if you should continue taking LAMICTAL XR.

2. Like other antiepileptic drugs, LAMICTAL XR may cause suicidal thoughts or actions in a very small number of people, about 1 in 500.

Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:

- thoughts about suicide or dying
- attempt to commit suicide
- new or worse depression
- new or worse anxiety
- feeling agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

Do not stop LAMICTAL XR without first talking to a healthcare provider.

- Stopping LAMICTAL XR suddenly can cause serious problems.
- Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

How can I watch for early symptoms of suicidal thoughts and actions?

- Pay attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled.
- Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

3. LAMICTAL XR may rarely cause aseptic meningitis, a serious inflammation of the protective membrane that covers the brain and spinal cord.

Call your healthcare provider right away if you have any of the following symptoms:

- Headache
- Fever
- Nausea
- Vomiting
- Stiff neck
- Rash
- Unusual sensitivity to light
- Muscle pains
- Chills
- Confusion
- Drowsiness

Meningitis has many causes other than LAMICTAL XR, which your doctor would check for if you developed meningitis while taking LAMICTAL XR.

LAMICTAL XR can have other serious side effects. For more information ask your healthcare provider or pharmacist. Tell your healthcare provider if you have any side effect that bothers you. Be sure to read the section below entitled “What are the possible side effects of LAMICTAL XR?”







4. Patients prescribed LAMICTAL have sometimes been given the wrong medicine because many medicines have names similar to LAMICTAL, so always check that you receive LAMICTAL XR.

Taking the wrong medication can cause serious health problems. When your healthcare provider gives you a prescription for LAMICTAL XR:

- Make sure you can read it clearly.
- Talk to your pharmacist to check that you are given the correct medicine.
- Each time you fill your prescription, check the tablets you receive against the pictures of the tablets below.

These pictures show the distinct wording, colors, and shapes of the tablets that help to identify the right strength of LAMICTAL XR. Immediately call your pharmacist if you receive a LAMICTAL XR tablet that does not look like one of the tablets shown below, as you may have received the wrong medication.

LAMICTAL XR (lamotrigine) Extended-Release Tablets

 25 mg, yellow with white center Imprinted with LAMICTAL XR 25	 50 mg, green with white center Imprinted with LAMICTAL XR 50	 100 mg, orange with white center Imprinted with LAMICTAL XR 100
 200 mg, blue with white center Imprinted with LAMICTAL XR 200	 250 mg, purple with white center Imprinted with LAMICTAL XR 250	 300 mg, gray with white center Imprinted with LAMICTAL XR 300

What is LAMICTAL XR?

LAMICTAL XR is a prescription medicine used:

- together with other medicines to treat primary generalized tonic-clonic seizures and partial onset seizures in people 13 years of age and older.
- alone to treat partial seizures when changing from certain other medicines used in people 13 years and older. It is not known if LAMICTAL XR is safe or effective in children less than 13 years of age. Other forms of lamotrigine can be used in children aged 2 to 12 years.

Who should not take LAMICTAL XR?

You should not take LAMICTAL XR if you have had an allergic reaction to lamotrigine or to any of the inactive ingredients in LAMICTAL XR. See the end of this leaflet for a complete list of ingredients in LAMICTAL XR.

What should I tell my healthcare provider before taking LAMICTAL XR?

Before taking LAMICTAL XR, tell your healthcare provider about all of your medical conditions, including if you:

- have had a rash or allergic reaction to another antiseizure medicine.
- have or have had depression, mood problems, or suicidal thoughts or behavior.
- are taking oral contraceptives (birth control pills) or other female hormonal medicines. Do not start or stop taking birth control pills or other female hormonal medicine until you have talked with your healthcare provider. Tell your healthcare provider if you have any changes in your menstrual pattern such as breakthrough bleeding. Stopping these medicines may

cause side effects (such as dizziness, lack of coordination, or double vision). Starting these medicines may lessen how well LAMICTAL XR works.

- are pregnant or plan to become pregnant. It is not known if LAMICTAL XR will harm your unborn baby. If you become pregnant while taking LAMICTAL XR, talk to your healthcare provider about registering with the North American Antiepileptic Drug Pregnancy Registry. You can enroll in this registry by calling 1-888-233-2334. The purpose of this registry is to collect information about the safety of antiepileptic drugs during pregnancy.
- are breastfeeding. LAMICTAL XR can pass into your breast milk. You and your healthcare provider should decide if you should take LAMICTAL XR or breastfeed.

Tell your healthcare provider about all the medicines you take or if you are planning to take a new medicine, including prescription and non-prescription medicines, vitamins, and herbal supplements. Using LAMICTAL XR with certain other medicines can affect each other, causing side effects.

How should I take LAMICTAL XR?

- Take LAMICTAL XR exactly as prescribed.
- Your healthcare provider may change your dose. Do not change your dose without talking to your healthcare provider.
- Do not stop taking LAMICTAL XR without talking to your healthcare provider. Stopping LAMICTAL XR suddenly may cause serious problems. For example, if you have epilepsy and you stop taking LAMICTAL XR suddenly, you may get seizures that do not stop. Talk with your healthcare provider about how to stop LAMICTAL XR slowly.
- If you miss a dose of LAMICTAL XR, take it as soon as you remember. If it is almost time for your next dose, just skip the missed dose. Take the next dose at your regular time. **Do not take 2 doses at the same time.**
- You may not feel the full effect of LAMICTAL XR for several weeks.
- If you have epilepsy, tell your healthcare provider if your seizures get worse or if you have any new types of seizures.
- LAMICTAL XR can be taken with or without food.
- Do not chew, crush, or divide LAMICTAL XR.
- Swallow LAMICTAL XR tablets whole.
- If you have trouble swallowing LAMICTAL XR Tablets, tell your healthcare provider because there may be another form of lamotrigine you can take.
- If you receive LAMICTAL XR in a blisterpack, examine the blisterpack before use. Do not use if blisters are torn, broken, or missing.

What should I avoid while taking LAMICTAL XR?

Do not drive a car or operate complex, hazardous machinery until you know how LAMICTAL XR affects you.

What are possible side effects of LAMICTAL XR?

- See “What is the most important information I should know about LAMICTAL XR?”

Common side effects of LAMICTAL XR include:

- Dizziness
- Tremor
- Double vision
- Nausea
- Vomiting
- Trouble with balance and coordination
- Anxiety

Other common side effects that have been reported with another form of lamotrigine include headache, sleepiness, blurred vision, runny nose, and rash.

Tell your healthcare provider about any side effect that bothers you or that does not go away. These are not all the possible side effects of LAMICTAL XR. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store LAMICTAL XR?

- Store LAMICTAL XR at room temperature between 59°F to 86°F (15°C to 30°C).
- **Keep LAMICTAL XR and all medicines out of the reach of children.**

General information about LAMICTAL XR

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use LAMICTAL XR for a condition for which it was not prescribed. Do not give LAMICTAL XR to other people, even if they have the same symptoms you have. It may harm them.

This Medication Guide summarizes the most important information about LAMICTAL XR. If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about LAMICTAL XR that is written for healthcare professionals.

For more information, go to www.lamictalxr.com or call 1-888-825-5249.

What are the ingredients in LAMICTAL XR?

Active ingredient: Lamotrigine.

Inactive ingredients: glycerol monostearate, hypromellose, lactose monohydrate, magnesium stearate, methacrylic acid copolymer dispersion, polyethylene glycol 400, polysorbate 80, silicon dioxide (25- and 50-mg tablets only), titanium dioxide, triethyl citrate, carmine (250-mg tablet only), iron oxide black (50-, 250-, and 300-mg tablets only), iron oxide yellow (25-, 50-, 100-mg tablets only), iron oxide red (100-mg tablet only), FD&C Blue No. 2 Aluminum Lake (200- and 250-mg tablets only). Tablets are printed with edible black ink.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

LAMICTAL XR is a trademark of GlaxoSmithKline.

DEPAKENE and DEPAKOTE are registered trademarks of Abbott Laboratories.



GlaxoSmithKline

Research Triangle Park, NC 27709

©2011, GlaxoSmithKline. All rights reserved.

June 2011

LXR:7MG